

# LEVAMISOLE



## POUR-ON FOR CATTLE

Levamisole (Clear) Anthelmintic  
Active Ingredient: 200g/L Levamisole base

### ADMINISTRATION

SHAKE WELL BEFORE USE.

VETMED Levamisole Pour-On contains Levamisole base, a member of the Levamisole family. It is effective against internal parasites sensitive to this family.

For accuracy, use an approved Pour-On gun. Check accuracy of Pour-On gun regularly. Apply slowly along animal's back according to live weight as shown in application table.

A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group (bulls, cows steers, calves etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on liveweight, to avoid excessive overdosing. Wet animals should not be treated. Do not use if rainfall is pending.

DOSE RATE: 1mL per 20kg bodyweight (10mg/kg Levamisole)			
WEIGHT RANGE	DOSE VOLUME	WEIGHT RANGE	DOSE VOLUME
50 – 75 kg	3.75 mL	351 – 400 kg	20.0 mL
75 – 125 kg	6.25 mL	401 – 450 kg	22.5 mL
126 – 150 kg	7.5 mL	451 – 500 kg	25.0 mL
151 – 200 kg	10.0 mL	501 – 550 kg	27.5 mL
201 – 250 kg	12.5 mL	551 – 600 kg	30.0 mL
251 – 300 kg	15.0 mL	601 – 650 kg	32.5 mL
301 – 350 kg	17.5 mL		

Cattle in excess of 500kg to be dosed at 1mL per 20kg bodyweight.

### WITHHOLD PERIODS

**MILK:** Milk intended for human consumption must be discarded during treatment and for no less than 24 hours following the last treatment.

**MEAT:** Animals producing meat and offal for human consumption must not be sold for slaughter either during treatment or within 10 days of cessation of the last treatment.

### COMBINED USE

Combined use of drench actives may improve efficacy and slow the development of resistance. Vetmed Levamisole Pour-On can be combined in equal volumes with Vetmed Abamectin Pour-On (A10786) to produce a stable mixture for Pour-On dosing of cattle. The mixture contains 5.0g/L Abamectin and 100g/L Levamisole. This combined solution must be well mixed before use and dosed at a rate of 1mL/10kg bodyweight. The mixture remains stable for up to 3 months if stored as per label conditions.

### STORAGE

Store in the original container, tightly closed and locked up in a safe place. Store below 25°C away from direct sunlight.

### PACK SIZES

VETMED Levamisole Pour-On is available in 5L and 2.5L packs.

VETMED Levamisole Pour-On is registered pursuant to the ACVM Act 1997 No. A10760.



09 274 8848 | [www.vetmed.co.nz](http://www.vetmed.co.nz)

Registered to: Nexan Corporation Limited, PO Box 217234, Botany Junction, Auckland

+1

LEVAMISOLE

BENEFITS

- Fast-acting
- Convenient Pour-On application
- 24hr milk withholding period
- Broad spectrum activity
- Effective against *Cooperia spp.*
- Effective against lungworm (Adult and immaturely *Dictyocaulus viviparus*)
- Solvent base delivering high blood levels of Levamisole
- Developed in New Zealand for New Zealand



CATTLE

FOR THE TREATMENT AND CONTROL OF  
GASTROINTESTINAL ROUNDWORMS AND  
LUNGWORM IN CATTLE.

Levamisole (Clear) Anthelmintic  
Active Ingredient: 200g/L Levamisole base