

**HANDLING PRECAUTIONS:**

Febantel possibly may affect development and the blood and haematopoietic system. Pyrantel possibly may affect the gastrointestinal system. Handle with care.

**FIRST AID:**

For advice contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately.

**DRONTAL® ALLWORMER FOR DOGS** is recommended for the control of all gastrointestinal worms in dogs including:

**Roundworm**

*Toxocara canis*, *Toxascaris leonina*

**Hookworm**

*Uncinaria stenocephala*, *Ancylostoma caninum*

**Whipworm**

*Trichuris vulpis*

**Tapeworm**

*Echinococcus granulosus* (Hydatid Tapeworm),  
*Taenia ovis* (sheep measles), *Dipylidium caninum*  
(common flea tapeworm), *Taenia hydatigena*,  
*Taenia pisiformis*

**DIRECTIONS FOR USE**

PUDDLES	Adults (over 6 months)	Pregnant/ Lactating Bitches
Treat at 2, 3, 4, 6, 9 and 12 weeks of age, then every 3 months	Every 3 months. Every 4 weeks in sheep measles infected locations	Prior to mating, 10 days before whelping and 2 and 4 weeks after whelping

DRONTAL ALLWORMER is given as a single dose by mouth and may be administered immediately before or with a meal. Fasting prior to dosing is NOT necessary. Weigh dogs before treatment. Treat all dogs and cats in the household.

**DOSAGE**

Bodyweight	Tablets
up to 5kg	½
5.1kg – 10kg	1
10.1kg – 20kg	2
20.1kg – 30kg	3
30.1 – 40kg	4

**RECOMMENDED TREATMENTS**

These recommendations are of a general nature. In specific instances where infestation is particularly heavy, or many dogs are housed together, a veterinary surgeon should be consulted. Flea tapeworms can reinfest dogs and begin producing segments 2-3 weeks after treatment. If segments reappear before the next DRONTAL treatment is due, either re-treat with DRONTAL ALLWORMER FOR DOGS or DRONCIT® as an intermediate, tapeworm only, treatment.

**WHY DO I NEED TO WORM MY DOG?**

Worms are extremely infectious. Puppies are born already infected with roundworms or infected with hookworms through the mother's milk, which may cause vomiting, diarrhoea, a 'pot belly' appearance and colic. Some adult worms can produce 20,000 eggs a day with some eggs surviving for up to five years on the ground, where they can be taken up by your pet by sniffing the ground, eating uncooked meats, rodents or swallowed during grooming. Worms can also be passed from animals to humans, with children particularly at risk. It is important that your dog receives proper worm treatment as worm burden symptoms are not always apparent.

**COMPATIBILITY**

DRONTAL can be used with other treatments, such as ADVANTAGE® for flea control.

**SAFETY**

DRONTAL offers gentle, effective control of worms. It may be given to all dogs including young puppies, old animals, greyhounds and is safe to use throughout pregnancy.

**STORAGE**

Store below 30°C (Room temperature)

DRONTAL ALLWORMER FOR DOGS is registered pursuant to the ACVM Act 1997 No. A5557. See [www.nzfsa.govt.nz/acvm](http://www.nzfsa.govt.nz/acvm) for registration conditions.



**Bayer**

Bayer New Zealand, Argus Place, Hillcrest, North Shore City, 0627. Phone: (09) 443 3093.

DRONTAL®, DRONCIT® and ADVANTAGE® are trademarks of Bayer AG Leverkusen. DRONCIT ACVM No. A2919. ADVANTAGE ACVM No. A7522.

Batch No:


Expiry:



# Drontal® ALLWORMER

## For Dogs

Controls all gastrointestinal worm species:  
Roundworm, Hookworm, Whipworm  
and Tapeworm



One tablet treats dogs  
up to 10kg.



Each tablet contains:  
50mg praziquantel,  
145mg pyrantel embonate  
and 150mg febantel.