

SLUG & SNAIL BAIT COLZACTIVE®

# **SNAIL AND SLUG BAIT**

For control of snails and slugs in agricultural, horticultural and ornamental crops and other situations where snails and slugs congregate

CONTAINS: 24.2 g/kg IRON PHOSPHATE ANHYDROUS (Equivalent to 30 g/kg IRON PHOSPHATE HYDRATED) in the form of a bait



Manufactured by and registered to: De Sangosse SA BP5I 47480 Pont Du Casse France

NET CONTENTS:

**15** 

Distributed by:
UPL New Zealand Limited
PO Box 51584, Pakuranga 2140
Auckland, New Zealand
Phone: 0800 100 325

Phone: 0800 100 325 www.upl-ltd.com/nz



## PLEASE READ THE LABEL COMPLETELY BEFORE USE

This product MUST NOT be used for any purpose or in any manner contrary to the controls and conditions of registration.



HSNO Classification: 6.4A, 9.1D

## WARNING

Causes eye irritation. Designed for biocidal action.

#### **PRECAUTIONS**

Keep out of reach of children. Wash hands after use.

#### STORAGE:

Store in the closed original container in a cool, well ventilated area, out of direct sunlight. When stored appropriately, this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of any product that is older than this.

#### DISPOSAL:

Dispose of this product by using in accordance with this label. Dispose of packaging to a suitable landfill.

#### FIRST AID:

If poisoning occurs, contact a doctor or the National Poisons Centre. Phone 0800 764 766.

#### ANIMAL SAFETY:

Low risk to pets and wildlife.

If your pet eats **IRONMAX Pro**, contact your veterinary surgeon immediately, as the consumption of large quantities may be toxic. The active ingredient in this product is iron phosphate.

# **GENERAL INFORMATION:**

**IRONMAX Pro** Snail and Slug Bait is a unique product containing iron phosphate, which is a natural component of soil, in a specially formulated bait. Any uneaten bait will degrade and become part of the soil.

After ingesting this bait (even in small amounts), snails and slugs cease feeding, which provides immediate protection to plants. Dead snails and slugs may not be visible as they often crawl away to secluded places to die. Plant protection will be observed in the dramatic decrease in plant damage.

Where the pest pressure is heavy, it may be necessary to repeat the application. Best results are achieved by application during mild damp weather when snails and slugs are most active. Snails need to be active to come into contact with the bait.

Scatter baits evenly using an appropriate applicator. DO NOT place in heaps. Clean up any spilled baits, including after filling applicator.

# TIMING:

Apply baits when crops are most vulnerable and when the pest is evident and active. For broadacre crops, start when rain/moisture triggers snail and slug activity in autumn or spring. Best results are achieved by applying just before or at crop emergence where crops are being sown. For transplants apply at time of transplanting. See your distributor for further advice.

## **DIRECTIONS FOR USE:**

| Crop/Situation   | Pest             | Rate      | Critical Comments   |
|--|------------------|-----------|---|
| Agricultural Crops Including Cereals, Oilseeds, Pulses, Pastures (established and seedling), Brassicas, Crop boundaries, Bare ground (including broadacre pre-emergent, pasture pre-establishment) | Snails and Slugs | 5-7 kg/ha | Broadcast evenly over<br>the surface of the<br>ground where snails<br>and slugs are active.<br>For most infestations,<br>use the lower rate.<br>For moderate to high<br>infestations, use a<br>higher rate. |
| Horticultural Crops<br>Including Vegetables,<br>Fruits (including<br>grapevines and citrus<br>orchards), Berry crops,<br>Ornamentals   |                  |           | Re-apply every 10-14<br>days or when pellets<br>have disappeared.   |
| Other Public authority and industrial areas, Roadsides, Turf and other situations where snails and slugs congregate  |                  |           | When applying to<br>crops, ensure pellets<br>do not become<br>lodged in plant<br>foliage.   |

**WITHHOLDING PERIOD:** All Edible Crops: Do not harvest for 1 day after application.

## **EMERGENCY RESPONSE:**

In a transport emergency dial 111, Police or Fire Brigade. For specialist advice in an emergency only, call 0800 CHEMCALL (0800 243 622).

# **SAFETY DATA SHEET:**

If additional safety information is required, refer to the Safety Data Sheet. For a copy visit www.upl-ltd.com/nz.

# **Exclusion of Liability: Please Read**

To the extent permitted by law, buyers and/or users of the goods accept that neither the manufacturer, UPL as distributor, nor any other distributor have any liability or responsibility whatsoever for any loss, damage or injury whether in contract or tort, whether direct, indirect or consequential howsoever arising in connection with the supply of these goods.

Registered pursuant to the ACVM Act 1997, No. P009801 See www.foodsafety.govt.nz for registration conditions. Approved pursuant to the HSNO Act 1996, Approval Code HSR000143 See www.epa.govt.nz for approval controls.

IRONMAX® is a registered trademark of DE SANGOSSE SAS

# UPL371/15KG

BATCH NO
DOM

24 Hour Emergency Response 0800 CHEM CALL 0800 243 622

